

IOPscience

Home Search Collections Journals About Contact us My IOPscience

Forthcoming (n, y) measurements on the Fe and Ni isotopes at CERN n_TOF

This article has been downloaded from IOPscience. Please scroll down to see the full text article. 2010 J. Phys.: Conf. Ser. 202 012026 (http://iopscience.iop.org/1742-6596/202/1/012026) View the table of contents for this issue, or go to the journal homepage for more

Download details: IP Address: 131.130.38.30 The article was downloaded on 06/05/2010 at 08:27

Please note that terms and conditions apply.

Forthcoming (n, γ) measurements on the Fe and Ni isotopes at CERN n_TOF

C. Domingo-Pardo^{1*}, S. Andriamonje⁴, J. Andrzejewski⁷, L. Audouin ²¹, V. Avrigeanu⁴⁰, F. Bečvář¹², F. Belloni³, B. Berthier²¹,

E. Berthoumieux⁴, M. Brugger¹⁷, M. Calviani²⁶, F. Calviño¹³,

D. Cano-Ott⁶, C. Carrapiço^{16,4}, F. Cerutti¹⁷, F. Chiaveri¹⁷, N. Colonna¹⁹, G. Cortes¹³, I. Dillmann⁴5, I. Duran⁵, A. Ferrari¹⁷,

W. Furman²⁴, S. Ganesan⁴¹, G. Giubrone², M.B. Gómez Hornillos¹³.

R. Gallino²⁵, I.F. Goncalves¹⁸, E. Gonzalez-Romero⁶, V. Gorlychev¹³,
F. Gramegna²⁶, C. Guerrero⁶, F. Gunsing⁴, S. Harissopulos³¹,

M. Heil¹, K. Ioannides⁸, E. Jericha¹⁰, F. Käppeler⁹, Y. Kadi¹⁷,

D. Karadimos⁸, M. Krtička¹², E. Lebbos¹⁷, C. Lederer⁴³, H. Leeb¹⁰, M. Lozano¹⁵, R. Losito¹⁷, J. Marganiec⁷, S. Marrone¹⁹, T. Martínez⁶,

C. Massimi³⁹, P. Mastinu²⁷, E. Mendoza⁶, A. Mengoni^{14,38},

P.M. Milazzo³, M. Mirea⁴⁰, M. Moinul¹⁹, M. Mosconi⁴⁴, R. Nolte⁴⁴, C. Paradela⁵, A. Pavlik³⁴, J. Perlowski²¹, R. Plag¹, J. Praena²⁶,

J.M. Quesada¹⁵, R. Reifarth¹, C. Rubbia²², R. Sarmento¹⁶,

G. Tagliente¹⁹, J.L. Tain², L. Tassan-Got²¹, D. Tarrio⁵, G. Vannini³⁹,

V. Variale¹⁹, P. Vaz¹⁶, A. Ventura³⁹, D. Villamarin⁶, V. Vlachoudis¹⁷, R. Vlastou³⁴, A. Wallner⁴³, and C. Wiess¹⁰

The n_TOF Collaboration

¹GSI Helmholtzzentrum für Schwerionenforschung GmbH, Darmstadt, Germany

²Instituto de Física Corpuscular, CSIC-Universidad de Valencia, Spain

³Istituto Nazionale di Fisica Nucleare, Trieste, Italy

⁴CEA/Saclay - DSM, Gif-sur-Yvette, France

⁵Universidade de Santiago de Compostela, Spain

 $^{6}\mathrm{Centro}$ de Investigaciones Energeticas Medioambientales y Technologicas, Madrid, Spain

⁷University of Lodz, Lodz, Poland

⁸University of Ioannina, Greece

⁹Forschungszentrum Karlsruhe GmbH (FZK), Institut für Kernphysik, Germany

¹0Atominstitut der Österreichischen Universitäten, Technische Universität Wien, Austria

¹¹Centre National de la Recherche Scientifique/IN2P3 - IReS, Strasbourg, France

¹²Charles University, Prague, Czech Republic

¹³Universitat Politecnica de Catalunya, Barcelona, Spain

¹⁴International Atomic Energy Agency - IAEA, Nuclear Data Section, Vienna, Austria ¹⁵Universidad de Sevilla, Spain

¹⁶Instituto Tecnológico e Nuclear(ITN), Lisbon, Portugal

 $^{17}\mathrm{CERN},$ Geneva, Switzerland

¹⁹Istituto Nazionale di Fisica Nucleare, Bari, Italy

²¹Centre National de la Recherche Scientifique/IN2P3 - IPN, Orsay, France

²⁴Joint Institute for Nuclear Research, Frank Laboratory of Neutron Physics, Dubna, Russia ²⁶Istituto Nazionale di Fisica Nucleare(INFN), Laboratori Nazionali di Legnaro, Italy

²⁷Centre National de la Recherche Scientifique/IN2P3 - CENBG, Bordeaux, France

²⁸Los Alamos National Laboratory, New Mexico, USA

²⁹Tokyo Institute of Technology, Tokyo, Japan

³⁰Oak Ridge National Laboratory, Physics Division, Oak Ridge, USA

- ³¹NCSR Demokritos, Athens, Greece
- ³³National Technical University of Athens, Greece
- ³⁴Institut für Isotopenforschung und Kernphysik, Universität Wien, Austria

³⁸ENEA, Bologna, Italy

³⁹Dipartimento di Fisica, Università di Bologna, and Sezione INFN di Bologna, Italy

⁴⁰National Institute of Physics and Nuclear Engineering - IFIN, Bucharest, Romania

⁴¹BARC, Mumbai, India ⁴²Argonne National Laboratory, Chicago, USA

⁴³VERA Laboratory - Isotopenforschung & Kernphysik, Faculty of Physics, University of Vienna, Austria

⁴⁴Physikalisch-Technische Bundesanstalt - PTB, Braunschweig, Germany

⁴⁵Physik-Department E12, Technische Universität München, Beschleunigerlaboratorium, Garching, Germany

E-mail: c.domingopardo@gsi.de

Abstract. An overview of the past, present and future research activities at the CERN neutron time of flight facility n_TOF is given, with special focus on the astrophysical aspects. During the first campaign (Phase I), neutron capture cross sections of relevance for several aspects of the s-process nucleosynthesis have been measured. A second campaign has recently started (Phase II), aiming at the study of the weak s-process component via accurate (n, γ) measurements of the Fe and Ni isotopes. Some changes in the facility will allow us to perform these measurements with improved experimental conditions.

1. Introduction

The n-TOF installation [1] aims at the measurement of accurate nuclear data for the Accelerator Driven System (ADS) project [2] and for nuclear astrophysics [3]. This facility combines a very large flight path of 185 m with an intense neutron spallation source, thus yielding an excellent energy resolution of 0.1% at 30 keV and a neutron intensity of more than 10^5 neutrons/pulse in the 1 keV-100 keV range. More details about the performance and commissioning of the installation can be found in Refs. [4, 5, 6]. In n-TOF Phase I, the first beam time was devoted to the further characterization of the resolution function [1], the ToF-neutron energy calibration [7], the spatial beam profile [8] and the detailed study of the pulse height weighting technique [9, 10]. A summary of the astrophysically relevant measurements carried out in Phase I, between 2002 and 2004, can be found in Sec. 2. Due to radiation-safety issues at the spallation source area, a new target has been constructed and the facility has resumed its activity in 2009. This upgrade and the new experimental programme are briefly described in Sec. 3.

2. n_TOF Phase I: measurements of astrophysical relevance for the s process

The list of isotopes measured in this phase at n_TOF is listed in Table 1. Particular emphasis deserves the measurement of a 151 Sm ($t_{1/2} = 93$ years) sample, with a radioactivity of 156 GBq, which was possible thanks to the high instantaneous neutron flux, low background and very small duty-cycle of n_TOF. Based on this experimental result, a detailed study of the thermal conditions during the recurrent thermal pulses of low mass AGB-stars could be performed [11, 12]. A detection system with reduced neutron sensitivity [13, 14] allowed the accurate measurement of low cross sections, i.e. the Mg isotopes and closed neutron shell nuclei (and neighboring isotopes) like Zr, La, Pb and Bi. The latters act as bottlenecks along the s-process path, hence accumulating large abundances.

Isotope	Astrophysically relevant aspects	References
^{24,25,26} Mg	Isotopic anomalies in meteorites	[15]
	Strength of ${}^{22}Ne(\alpha, n)$ source	
^{90,91,92,93,94,96} Zr	Weak/Main s process components	[16, 17, 18]
	Bottleneck at $N = 50$	
	⁹⁵ Zr branching sensitive to neutron flux	
¹³⁹ La	Indicator of s-process activity	[19]
	Bottleneck at $N = 82$	
151 Sm	Branching of s-process path	[11, 20, 12]
	Thermal conditions of AGB stars	
$^{186,187,188}\mathrm{Os}$	Nuclear cosmo-chronology	[21]
^{204,206,207} Pb, ²⁰⁹ Bi	Termination of s-process path	[22, 23, 24, 25]
	Constraint for r-process model calculations	
	Bottleneck at $N = 126$	
¹³⁹ La ¹⁵¹ Sm ^{186,187,188} Os ^{204,206,207} Pb, ²⁰⁹ Bi	Bottleneck at $N = 50^{95}$ Zr branching sensitive to neutron flux Indicator of s-process activity Bottleneck at $N = 82$ Branching of s-process path Thermal conditions of AGB stars Nuclear cosmo-chronology Termination of s-process path Constraint for r-process model calculations Bottleneck at $N = 126$	[19] $[11, 20, 12]$ $[21]$ $[22, 23, 24, 25]$

Table 1. (n, γ) measurements made during n_TOF Phase I.

3. n_TOF Phase II: s process in massive stars

Some aspects have been improved in the facility, particularly concerning the spallation source. On one hand, the target is equipped now with the possibility to include a borated or heavy water moderation layer, which will allow one to highly reduce the γ -rays produced by neutron captures in the hydrogen of the moderator, thus improving the peak-to-background ratio in the keV-MeV region by more than a factor of 5. On the other hand, the new spallation target has been designed in such a way, that it may serve in the future as a common neutron source for a second experimental area at a distance of 20 m. The latter would provide an increase in sensitivity by two to three orders of magnitude.

Since May 2009 a series of measurements have been carried out for commissioning the upgraded facility. The neutron flux has been determined from measurements with a micromegas detector containing ¹⁰B and ²³⁵U deposits, from the ⁶Li(n, α) reaction measured with a silicon array (SiMon) surrounding a thin Li-foil, and from the ²³⁵U(n,f) measured with a fission chamber from PTB (Physikalisch-Technische Bundesanstalt) Braunschweig. An absolute value of the neutron fluence at 4.9 eV was obtained from the ¹⁹⁷Au(n, γ) measurement with the Total Absorption Calorimeter [26]. A 2D-measurement of the spatial beam profile versus the neutron energy has been carried out using micromegas detectors. The background conditions in the experimental area will be characterized from measurements made with lead, graphite and empty samples. Finally, the resolution function will be characterized from the measurement of a ⁵⁶Fe sample.

Table 2. (n, γ) measurements planned for n_TOF Phase II.

Isotope	Astrophysically relevant aspects
^{54,56,57,58} Fe	Prime s-process contributions to the galactic chemical
^{58,60,61,62,(63),64} Ni	enrichment and quantitative study of the weak s process

Nuclear Physics in Astrophysics IV	IOP Publishing
------------------------------------	----------------

doi:10.1088/1742-6596/202/1/012026

The weak s process has been found to be extremely sensitive to the neutron capture cross sections of the isotopes near the seed distribution around Fe [27, 28]. Therefore, accurate (n, γ) measurements on these nuclei will be the main goal of the present campaign (see Table 2). Indeed, the neutron exposure during the weak s process is too small to achieve flow equilibrium, which means that the cross section uncertainty on one isotope affects not only its own abundance, but also the production of the subsequent higher mass isotopes. Due to this propagation effect, it is important to measure the cross section of all involved Fe-group nuclei over a broad energy range and with high accuracy [29].

References

- U Abbondanno et al 2002 (Preprint CERN n_TOF Facility: Performance Report, CERN/INTC-0-011 INTC-2002-037 CERN-SL-2002-053ECT)
- [2] C Rubbia et al 2000 (Preprint 1st Management Report of the n_TOFND-ADS EC programme under contract no. FIKW-CT-2000-00107)
- [3] Mengoni A 2002 Astrophysics, Symmetries, and Applied Physics at Spallation Neutron Sources ed PE Koehler et al p 25
- [4] C Borcea et al 2003 Nuclear Instruments and Methods in Physics Research A 513 524–537
- [5] U Abbondanno et al 2001 (Preprint Tech. Rep. CERN-INTC-2001-038)
- [6] A Ferrari et al 2002 (Preprint Tech. Rep. CERN-SL-2002-011 ECT)
- [7] G Lorusso et al 2004 Nucl. Instr. and Meth. Phys. Res. A 532 622-630
- [8] J Pancin et al 2004 Nuclear Instruments and Methods in Physics Research A 524 102–114
- [9] JL Tain et al 2002 Jour. of Nucl. Sci. and Tech. vol 2 (Tsukuba) p 689
- [10] U Abbondanno et al 2004 Nuclear Instruments and Methods in Physics Research A 521 454-467
- [11] U Abbondanno et al 2004 Phys. Rev. Lett. 93 161103
- [12] S Marrone et al 2006 Phys. Rev. C 73 034604
- [13] R Plag et al 2003 Nuclear Instruments and Methods in Physics Research A 496 425-436
- [14] S Marrone et al 2004 Nuclear Instruments and Methods in Physics Research A 517 389–398
- [15] E Strandberg 2007 AGB Star nucleosynthesis of the Magnesium Isotopes Ph.D. thesis University of Notre Dame http://physics.nd.edu/Pdf/Theses/StrandbergE042007.pdf
- [16] G Tagliente et al 2006 International Symposium on Nuclear Astrophysics Nuclei in the Cosmos
- $\left[17\right]$ G Tagliente et al 2008 Phys. Rev. C $\mathbf{77}$ 035802
- [18] G Tagliente et al 2008 Phys. Rev. C ${\bf 78}$ 045804
- [19] R Terlizzi et al 2007 Phys. Rev. C 75 035807 (Preprint arXiv:nucl-ex/0610034)
- [20] S Marrone et al 2005 Nucl. Phys. A 758 533–536
- [21] M Mosconi et al 2007 Prog. Part. and Nucl. Phys. 59 165–173
- [22] C Domingo-Pardo et al 2006 Phys. Rev. C 74 025807
- [23] C Domingo-Pardo et al 2006 Phys. Rev. C 74 055802
- [24] C Domingo-Pardo et al 2007 Phys. Rev. C 75 015806 (Preprint arXiv:nucl-ex/0610033)
- [25] C Domingo-Pardo et al 2007 Phys. Rev. C 76 045805 (Preprint arXiv:nucl-ex/0707.3679)
- [26] C Guerrero et al 2009 American Institute of Physics Conference Series (American Institute of Physics Conference Series vol 1090) ed Jolie J, Zilges A, Warr N and Blazhev A pp 372–375
- [27] CM Raiteri et al 1991 The Astr. Jour. **371** 665–672
- [28] T Rauscher et al 2002 The Astr. Jour. 576 323-348 (Preprint arXiv:astro-ph/0112478)
- [29] M Heil et al 2008 Phys. Rev. C 77 015808