

The Production of ^{53}Mn in a Fusion-Environment

C. Lederer^a, A. Wallner^a, I. Dillmann^b, T. Faestermann^b, A. Klix^c,
G. Korschinek^b, J. Lachner^b, M. Poutivtsev^b, G. Rugel^b, K. Seidel^c, and H. Vonach^a


^aIsotopenforschung und Kernphysik, Fakultät für Physik, Universität Wien

^bPhysik Department, Technische Universität München

^cInst. f. Kern- und Teilchenphysik, TU Dresden und Forschungszentrum Dresden-Rossendorf

Motivation

Reaction:



http://ec.europa.eu/commission_barroso/potocnik/images/dossier/iter.jpg


Claudia Lederer




universität
wien

Motivation

Reaction:




A. Wallner, 2000



http://ec.europa.eu/commission_barroso/potocnik/images/dossier/iter.jpg

Motivation

Reaction:




A. Wallner, 2000

Isotope	Half-life (y)	Reaction
^{14}C	5730	$^{14}\text{N}(n,p)$
^{26}Al	$7.2 \cdot 10^5$	$^{27}\text{Al}(n,2n)$
^{53}Mn	$3.7 \cdot 10^6$	$^{54}\text{Fe}(n,2n)$
		$^{54}\text{Fe}(n,np+d)$
^{55}Fe	2.73	$^{56}\text{Fe}(n,2n)$
^{63}Ni	100	$^{63}\text{Cu}(n,p)$
		$^{64}\text{Ni}(n,2n)$
^{59}Ni	$7.6 \cdot 10^4$	$^{60}\text{Ni}(n,2n)$
^{93}Zr	$1.5 \cdot 10^6$	$^{94}\text{Zr}(n,2n)$
		$^{93}\text{Nb}(n,p)$
^{91}Nb	680	$^{92}\text{Nb}(n,2n)$
^{92}Nb	$3.2 \cdot 10^7$	$^{93}\text{Nb}(n,2n)$
$^{93\text{m}}\text{Nb}$	16.1	$^{94}\text{Mo}(n,x)$
^{94}Nb	$2.0 \cdot 10^4$	$^{94}\text{Mo}(n,p)$
^{93}Mo	3500	$^{94}\text{Mo}(n,2n)$

Claudia Lederer

Motivation

Reaction:




A. Wallner, 2000

Isotope	Half-life (y)	Reaction
^{14}C	5730	$^{14}\text{N}(n,p)$
^{26}Al	$7.2 \cdot 10^5$	$^{27}\text{Al}(n,2n)$
^{53}Mn	$3.7 \cdot 10^6$	$^{54}\text{Fe}(n,2n)$
		$^{54}\text{Fe}(n,np+d)$
^{55}Fe	2.73	$^{56}\text{Fe}(n,2n)$
^{63}Ni	100	$^{63}\text{Cu}(n,p)$
		$^{64}\text{Ni}(n,2n)$
^{59}Ni	$7.6 \cdot 10^4$	$^{60}\text{Ni}(n,2n)$
^{93}Zr	$1.5 \cdot 10^6$	$^{94}\text{Zr}(n,2n)$
		$^{93}\text{Nb}(n,p)$
^{91}Nb	680	$^{92}\text{Nb}(n,2n)$
^{92}Nb	$3.2 \cdot 10^7$	$^{93}\text{Nb}(n,2n)$
$^{93\text{m}}\text{Nb}$	16.1	$^{94}\text{Mo}(n,x)$
^{94}Nb	$2.0 \cdot 10^4$	$^{94}\text{Mo}(n,p)$
^{93}Mo	3500	$^{94}\text{Mo}(n,2n)$

Claudia Lederer

Available data for $^{54}\text{Fe}(\text{n},\text{np+d})^{53}\text{Mn}$ cross-sections



<http://www-nds.iaea.org/>

Cross-section determination

Cross-section determination

- Irradiation of highly enriched ^{54}Fe with quasi-monoenergetic neutrons from 13-15 MeV at FZD

Cross-section determination

- Irradiation of highly enriched ^{54}Fe with quasi-monoenergetic neutrons from 13-15 MeV at FZD
- Adding stable ^{55}Mn as spike

Cross-section determination

- Irradiation of highly enriched ^{54}Fe with quasi-monoenergetic neutrons from 13-15 MeV at FZD
- Adding stable ^{55}Mn as spike
- Chemical removal of Fe and other impurities from sample

Cross-section determination

- Irradiation of highly enriched ^{54}Fe with quasi-monoenergetic neutrons from 13-15 MeV at FZD
- Adding stable ^{55}Mn as spike
- Chemical removal of Fe and other impurities from sample
- Measurement of ^{53}Mn / ^{55}Mn with AMS at MLL Munich



Cross-section determination

- Irradiation of highly enriched ^{54}Fe with quasi-monoenergetic neutrons from 13-15 MeV at FZD
- Adding stable ^{55}Mn as spike
- Chemical removal of Fe and other impurities from sample
- Measurement of ^{53}Mn / ^{55}Mn with AMS at MLL Munich

$$N_{^{53}\text{Mn}} = N_{^{54}\text{Fe}} \cdot \phi \cdot \sigma(E_n) \cdot f$$



Cross-section determination

- Irradiation of highly enriched ^{54}Fe with quasi-monoenergetic neutrons from 13-15 MeV at FZD
- Adding stable ^{55}Mn as spike
- Chemical removal of Fe and other impurities from sample
- Measurement of ^{53}Mn / ^{55}Mn with AMS at MLL Munich

$$N_{^{53}\text{Mn}} = N_{^{54}\text{Fe}} \cdot \phi \cdot \sigma(E_n) \cdot f$$

$$\rightarrow \sigma(E_n) = \frac{N_{^{53}\text{Mn}}}{N_{^{55}\text{Mn}}} \frac{N_{^{55}\text{Mn}}}{N_{^{54}\text{Fe}}} \frac{1}{\phi \cdot f}$$



Cross-section determination

- Irradiation of highly enriched ^{54}Fe with quasi-monoenergetic neutrons from 13-15 MeV at FZD
- Adding stable ^{55}Mn as spike
- Chemical removal of Fe and other impurities from sample
- Measurement of $^{53}\text{Mn} / ^{55}\text{Mn}$ with AMS at MLL Munich

$$N_{^{53}\text{Mn}} = N_{^{54}\text{Fe}} \cdot \phi \cdot \sigma(E_n) \cdot f$$

$$\rightarrow \sigma(E_n) = \frac{N_{^{53}\text{Mn}}}{N_{^{55}\text{Mn}}} \frac{N_{^{55}\text{Mn}}}{N_{^{54}\text{Fe}}} \frac{1}{\phi \cdot f}$$

AMS



Cross-section determination

- Irradiation of highly enriched ^{54}Fe with quasi-monoenergetic neutrons from 13-15 MeV at FZD
 - Adding stable ^{55}Mn as spike
 - Chemical removal of Fe and other impurities from sample
 - Measurement of ^{53}Mn / ^{55}Mn with AMS at MLL Munich

$$N_{^{53}\text{Mn}} = N_{^{54}\text{Fe}} \cdot \phi \cdot \sigma(E_n) \cdot f$$

$\rightarrow \sigma(E_n) = \frac{N_{^{53}\text{Mn}}}{N_{^{55}\text{Mn}}} \cdot \frac{N_{^{55}\text{Mn}}}{N_{^{54}\text{Fe}}} \cdot \frac{1}{\phi \cdot f}$

AMS

mass measurement

Cross-section determination

- Irradiation of highly enriched ^{54}Fe with quasi-monoenergetic neutrons from 13-15 MeV at FZD
 - Adding stable ^{55}Mn as spike
 - Chemical removal of Fe and other impurities from sample
 - Measurement of ^{53}Mn / ^{55}Mn with AMS at MLL Munich

$$N_{^{53}\text{Mn}} = N_{^{54}\text{Fe}} \cdot \phi \cdot \sigma(E_n) \cdot f$$

$$\rightarrow \sigma(E_n) = \frac{N_{^{53}\text{Mn}}}{N_{^{55}\text{Mn}}} \frac{N_{^{55}\text{Mn}}}{N_{^{54}\text{Fe}}} \frac{1}{\phi \cdot f}$$

AMS

mass measurement

AMS


mass measurement

monitor reactions

Neutron Irradiations


Neutron Irradiations

- Neutron generator at Forschungszentrum Dresden: $t(d,n)^4\text{He}$
- 11 positions with ^{13}C , ^{14}N , ^{54}Fe , $^{\text{nat}}\text{Fe}$ samples
- 4 positions for $^{54}\text{Fe}(n,np+d) ^{53}\text{Mn}$




Neutron Irradiations

- Neutron generator at Forschungszentrum Dresden: $t(d,n)^4\text{He}$
- 11 positions with ^{13}C , ^{14}N , ^{54}Fe , $^{\text{nat}}\text{Fe}$ samples
- 4 positions for $^{54}\text{Fe}(n,np+d) ^{53}\text{Mn}$





A. Wallner, 2000



Neutron Irradiations

- Neutron generator at Forschungszentrum Dresden: $t(d,n)^4\text{He}$
- 11 positions with ^{13}C , ^{14}N , ^{54}Fe , $^{\text{nat}}\text{Fe}$ samples
- 4 positions for $^{54}\text{Fe}(n,np+d) ^{53}\text{Mn}$






$^{53}\text{Mn}/^{55}\text{Mn}$ measurements with AMS (AMS....Accelerator Mass Spectrometry)




$^{53}\text{Mn}/^{55}\text{Mn}$ measurements with AMS (AMS....Accelerator Mass Spectrometry)



Munich 14 MV tandem accelerator

$^{53}\text{Mn}/^{55}\text{Mn}$ measurements with AMS (AMS....Accelerator Mass Spectrometry)



^{53}Mn - stable isobar ^{53}Cr


Munich 14 MV tandem accelerator

Isobar suppression: GAMS

- GAMS.... Gas-filled Analyzing Magnet System


Isobar suppression: GAMS

- GAMS.... Gas-filled Analyzing Magnet System
- Principle:



Isobar suppression: GAMS

- GAMS.... Gas-filled Analyzing Magnet System
- Principle:




Cr-suppression:


- Magnet: 10^3
- Ionization chamber: 10^6

[fwd](#)


Results



Results




Results






Thank you!



Reference Material




Reference Material




Cr-blank: ^{55}Mn with 1000 ppm ^{53}Cr 


Cr-blank: ^{55}Mn with 1000 ppm ^{53}Cr 

Cr-blank: ^{55}Mn with 1000 ppm ^{53}Cr 


Cr-blank: ^{55}Mn with 1000 ppm ^{53}Cr



Cr-blank: ^{55}Mn with 1000 ppm ^{53}Cr



Cr-blank: ^{55}Mn with 1000 ppm ^{53}Cr



Cr-blank: ^{55}Mn with 1000 ppm ^{53}Cr

